

**EXTRACTION OF VOLATILE OIL FROM GINGER (*Zingiber officinale*
Roscoe)**

WAN MOHD FATHULMUIN BIN WAN HASSAN

**Final Year Project Report Submitted in
Partial Fulfilment of the Requirements for the
Degree of Bachelor of Science (Hons.) Chemistry
in the Faculty of Applied Sciences
Universiti Teknologi MARA**

JANUARY 2017

ABSTRACT

Ginger (*Zingiber officinale* Roscoe) is well known as a spice which has been used over 2000 year ago. It can be found in tropical region such as Malaysia, Indonesia and Thailand. Ginger has highly used in the worldwide as a spice, herbal remedy, and flavoring agent etc. This final year project is about the Extraction of Volatile Oil from ginger (*Zingiber officinale* Roscoe) that is obtained from Malaysia local market. Hydrodistillation process was conducted in order to extract the essential oil from the rhizome ginger. It was separate using hexane and chloroform in order to determine the desire compound at different polarity of solvent. The major compound that will present in the essential oil of ginger is depending on the extraction time of plant studied. The percent yield obtained from hexane extract for two hour and four hour of extraction was 0.1059% and 0.1663%. Meanwhile, the percent yield from choloroform extract for two hour and four hour of extraction was 0.0591% and 0.1358%. GC-MS analysis was used in order to determine desire compound in the ginger oil. Seven chemical compounds was founds on chloroform extract at two and four hour of extraction time which is 2-hydroxycineol (2 hr : 71.6%), 8-hydroxyneomenthol (2 hr : 31.3%), oleic acid (2 hr : 29.8%), 3-deoxyestradiol (2 hr : 56.8%) and 6-(1-hydroxy-1-methylethyl)-3-methyl-2-cyclohexenol (4 hr : 26.0%), adipic acid, bis(2-ethylhexyl) ester (4 hr : 20.8%) and 2-methylenecholestan-3-ol (4 hr : 22.5%), respectively. Meanwhile, nine chemical compounds was found on hexane extract at two and four hour of extraction time, β -Linalool (2 hr : 70.2% ; 4 hr : 61.7%), α -terpineol (2 hr : 58.3%), β -citral (2 hr

: 72.6% ; 4 hr : 46.1%), α -citral (2 hr : 73.8%), α -curcumene (2 hr : 87.2% ; 4 hr : 76.6%), zingiberenol (2 hr : 56.2%) and 6-methyl-5-heptene-2-one (4 hr : 77.3%), nerolidol (4 hr : 45.2%), retinoic acid, methyl ester (4 hr : 52.9%). Nine compounds, 2-hydroxycineol, 8-hydroxyneomenthol, oleic acid, 3-deoxyestradiol 6-(1-hydroxy-1-methylethyl)-3-methyl-2-cyclohexenol, adipic acid, bis(2-ethylhexyl) ester and 6-methyl-5-heptene-2-one and retinoic acid, methyl ester was reported for a first time in Zingiberaceae family while 2-methylenecholestan-3-ol was first time found in this species.

TABLE OF CONTENTS

	PAGE
ACKNOWLEDGEMENT	ii
TABLE OF CONTENTS	iii
LIST OF TABLES	v
LIST OF FIGURES	vi
LIST OF ABBREVIATIONS	vii
ABSTRACT	ix
ABSTRAK	xi

CHAPTER 1 INTRODUCTION

1.1	Background of Study	1
1.2	Botany of plant	2
	1.2.1 Family of <i>Zingiber officinale</i> Roscoe species	2-3
1.3	Problem Statement	4
1.4	Significance of Study	4
1.5	Objectives of Study	5
1.6	Scope of Study	5

CHAPTER 2 LITERATURE REVIEW

2.1	Hydrodistillation and secondary metabolites of medicinal plant	6
2.2	Essential oils from fresh ginger (<i>Zingiber officinale</i> Roscoe)	6-10
2.3	Anti-microbial activity of <i>Zingiber officinale</i> Roscoe	11
2.4	Anti-cancer properties of <i>Zingiber officinale</i> Roscoe	12
2.5	Anti-Oxidant Activities <i>Zingiber officinale</i> Roscoe	13

CHAPTER 3 METHODOLOGY

3.1	Plant Materials	14
3.2	Instruments	14
	3.2.1 Gas Chromatography Mass Spectra	14
3.3	Extraction and analysis of chemical composition from fresh rhizome of <i>Zingiber officinale</i>	15
3.4	Gas Chromatography Mass Spectrometry Analysis	16
3.5	Flow of methodology	17

CHAPTER 4 RESULT AND DISCUSSION

4.1	Hydrodistillation extraction of rhizome of <i>Zingiber officinale</i>	18-20
4.2	GC-MS analysis	21-22
	4.2.1 Chemical structure in ginger oil using hexane extract	23-25
	4.2.2 Chemical structure in ginger oil using chloroform extract	26-31

CHAPTER 5 CONCLUSION AND RECOMMENDATION 32-33

REFERENCES 34-36

CURICULUM VITAE 37-38